

How to do Smooth Convergence Exercises:

This leaflet is for patients who have been diagnosed with Convergence weakness

These exercises are a very effective treatment when they are done properly.

When you were in clinic your Orthoptist will have given you a suitable target to look at when performing these exercises. She will have also explained how to perform the exercises correctly. However, this is often a lot to take in, so this simple method should help you to do your exercises correctly.

Method:

1. If you require **reading glasses these should be worn** when doing these exercises. If the Orthoptist has put a Fresnel prism over one lens of your reading glasses this should be kept on.
2. Take your target and hold it at arm's length.
3. Keep both eyes open at all times when looking at your target.
4. If you can see only one of the target, **very slowly** move it towards your nose until you see two.
5. Stop and hold your arm still when you see two of the target. Holding the target steady, stare at it and do **NOT** look away for at least 15 seconds. Try to make and keep the target one.
6. If you can make and keep the target one then you can very slowly move the target further in towards your nose.
7. Stop and hold your arm exactly where it is when you see two of the target. Keeping both eyes open at all times, stare at the target for at least 15 seconds. Try to make the target one.
8. If you cannot make the target one, very slowly move it away from your nose until it is single. Then try to move it forwards keeping it single aiming to get as close to your nose as possible without seeing two of the target.
9. Keep doing this for 2 to 3 minutes aiming to keep the target single at all times.
10. You should aim to do your exercises 4 to 5 times a day for no longer than 2 to 3 minutes at a time.
11. After doing these exercises it is important to look into the distance to relax your eyes.



Simple Tips:

- Keep both eyes open all the time.
- Always wear your reading glasses.
- Don't rush, take your time – move the target in very slowly towards your nose so that you accurately recognise when you are seeing two of the target.
- Be patient – you will not be able to keep the target one and get to your nose immediately. It will take time and practice.

What to do if your target is not single at arm's length:

Rest your target against a wall or stick it to the wall so that it is on eye level. Step away from the target until it is single. Then move towards the target very slowly until it goes double. When it goes into two, stand very still and stare at the target for at least 15 seconds. If you can make it single move very slowly further in towards the target, trying to keep it single. If you cannot make it single step back until it becomes one then try to keep it single and move forwards again. Do this for 4 minutes at a time and do 4 to 5 sessions a day.

If you have any questions about your treatment, please ask your Orthoptist or contact us on the phone numbers below:

Orthoptic Departments

Pilgrim Hospital, Boston
Orthoptic Department: 01205 446474

Lincoln County Hospital
Orthoptic Department: 01522 573378

Monday to Friday 9.00am to 5.00pm

www.ulh.nhs.uk

References

If you require a full list of references for this leaflet please email patient.information@ulh.nhs.uk

The Trust endeavours to ensure that the information given here is accurate and impartial.

If you require this information in another language, large print, audio (CD or tape) or braille, please email the Patient Information team at patient.information@ulh.nhs.uk