

What happens after surgery?

Once the operation is finished you will be moved to an area called “recovery” within the theatre complex. You will stay there for at least 4 hours when your blood pressure and neurological signs will be closely monitored. Once the recovery staff are satisfied you are stable you can have something to eat and drink. After this you will return to the surgical ward. Most patients spend 1 to 2 days in hospital and only require simple painkillers.

Risks

Local anaesthetics are generally very safe and serious problems are rare. If you develop a stroke during the operation the anaesthetist might convert to a general anaesthetic to continue with the operation safely. If you have any questions regarding the anaesthetic please discuss with the pre op nurse or anaesthetist who will be happy to help.

Consultant Vascular Anaesthetists are:

Dr W Butt
Dr S Panjwani
Dr S Joachim
Dr R Jaganathan
Dr M Chablani

References

If you require a full list of references for this leaflet please email patient.information@ulh.nhs.uk

The Trust endeavours to ensure that the information given here is accurate and impartial.



If you require this information in another language, large print, audio (CD or tape) or braille please email the Patient Information team at patient.information@ulh.nhs.uk



Local Anaesthesia for Carotid Endarterectomy

Department of Anaesthesia
Pilgrim Hospital
01205 445293
www.ulh.nhs.uk

Aim of the leaflet

This leaflet is aimed at patients having a carotid endarterectomy surgery under local anaesthesia.

This leaflet aims to tell you why you need a local anaesthetic and what will happen before, during and after the operation.

Why do I need surgery?

You have narrowing of the artery that supplies blood to your brain (carotid artery). Either you have had a “mini stroke” (TIA) or you do not have any symptoms. The danger is that you could have a major stroke. The aim of this surgery is to decrease the chance of you having another mini stroke or major stroke.

Surgery for clearing a narrowing or blockage in the carotid artery is a major procedure. For most patients the safest technique is a local anaesthetic, which involves numbing the side of the neck being operated on. This allows the doctors looking after you to monitor you in the best possible way.

Why do I have a local anaesthetic?

During the surgery, a clamp is placed in the carotid artery which may reduce the blood supply to your brain. The best way of assessing this effect is to have you awake and talking to us. If you develop any symptoms similar to a “mini stroke” the doctors can modify your treatment accordingly.

What happens before the operation?

The anaesthetist will come to see you the night before or on the day of your operation. You will have a local anaesthetic cream and a dressing applied to your neck. You will be encouraged to empty your bladder before going to theatre as it avoids the need for a urinary catheter.

What happens in the anaesthetic room?

Once you are in the anaesthetic room you will be attached to some routine monitoring devices. (ECG, Blood pressure cuff, Oxygen monitor in your finger). The anaesthetist will put two drips in your arm that is on the same side as your operation. He will also give you some medication to help you relax. He will clean your neck with an antiseptic solution and then make some marks in your neck. He will then give you 3 to 5 injections in your neck, always explaining what he is doing.

What happens in theatre?

In the operating theatre there will be a full team of staff attending to you. The surgeon will clean your neck and place a plastic drape over your neck. This is the surgical field and helps to keep everything clean and stops the wound from getting infected. You will not be able to see the operating site. You will have a member of staff at your bedside who will keep you company.

You will be given a rubber ball in your other hand (the one without any drips in it) and will be asked to squeeze the ball during the operation. The anaesthetist will be in theatre throughout the procedure keeping a close eye on your vital signs.

If you feel any pain during the operation the surgeon can give you more local anaesthetic to make it better. It is important that you don't make any movements with your hand. You can communicate any concerns to the nurse sat next to you or the anaesthetist.

The procedure takes 2 to 3 hours to complete, you will be able to talk to people during the operation but there will be times when you have to be very still and not talking. The theatre team will do everything they can to keep you as comfortable as possible during the operation. Your co-operation during the procedure is very important.