

HDR Brachytherapy (Internal Radiotherapy) Endometrium

Radiotherapy Department
Lincoln County Hospital
www.ulh.nhs.uk

Gynaecological Brachytherapy (Internal Radiotherapy) Using microSelectron HDR

The aim of this leaflet is to explain what brachytherapy is and the treatment procedure.

This type of radiotherapy treatment is used for treating gynaecological cancers. It is routinely carried out as an out patient procedure.

What is microSelectron HDR?

This is the name of the machine that delivers the internal radiotherapy. It uses a small radioactive pellet that passes into thin, hollow tubes called applicators. The radioactive pellet does not touch you. You will not be radioactive following the treatment.

What will happen?

The preparation and treatment are carried out in the same room, within the brachytherapy suite. Family and/or friends will not be able to enter this area.

You have had a hysterectomy (womb removal) and do not need an anaesthetic for the procedure. For the first treatment the doctor or radiographer will carry out an internal examination using a speculum, but this is not required for the subsequent session.

The applicator will be inserted once the treatment time has been calculated. This is calculated before the treatment and takes approximately 10 minutes. The applicator is held in place by a support bracket on the bed. There may be very slight discomfort for a short time.

What happens during treatment?

You are not required to have a full bladder for this treatment.

You will remain in the same position until the end of the treatment session. This will be lying on your back with legs flat.

You will have pillows under your head and support sponges under your ankles (and knees if required).

A radiographer will be with you until the treatment is switched on.

There can be no members of staff in the treatment room during treatment but you will be watched continually on a closed circuit TV.

How long is the treatment?

The treatment time is usually less than 10 minutes for each session. You will be told the treatment time before it commences.

How many treatments will I have?

The treatment is usually given twice, one per week and one week apart, usually on a Thursday. The doctor will explain to you how many treatments you need.

What happens when the treatment is finished?

A radiographer will remove the applicators and you can then get dressed and leave the department when you feel ready. If you are an in patient a porter will return you to the ward.

Short term problems you may have following treatment

You may experience looser bowels for a few days after each treatment. You may also feel the desire to pass urine more often.

Vaginal Dilation After Radiotherapy Treatment

This information is provided in addition to any information given to you by your consultant.

Radiotherapy to the vaginal area can cause the vagina to become shorter, narrower and less elastic than before treatment, as well as being drier due to loss of vaginal secretions.

Vaginal dilation will reduce the amount of scar tissue formation by gently stretching the tissues, helping to keep the vagina more supple and future medical examinations more comfortable. Dilation can be achieved by using a dilator, sexual intercourse, or a combination of the two.

A radiographer will discuss the use of dilators after you have been given your last brachytherapy treatment and will provide you with a dilator pack. A vaginal dilator is a smooth plastic tube and your dilator pack will contain 4 different sizes.

Useful websites:

www.jostrust.org.uk
www.cancerresearch.org.uk
www.macmillan.org.uk
www.daisynetwork.org.uk
www.infertilitynetwork.uk
www.maggiescentres.org
www.mariecurie.org.uk
www.nhsdirect.nhs.uk
www.eastmidlandscancernetwork.nhs.uk

Telephone Numbers

If you have any queries regarding your brachytherapy treatment contact the relevant gynaecology specialist nurse or brachytherapy specialist.

Gynaecology nurses: Lincoln 01522 573126
Boston 01205 445431

Lucy Darby/Melanie Fisher:
Advanced Practitioners Brachytherapy 01522 572243

Dr M Panades via his secretary: Lincoln 01522 572218
Boston 01205 445262/446432

References

If you require a full list of references for this leaflet please email patient.information@ulh.nhs.uk

The Trust endeavours to ensure that the information given here is accurate and impartial.



If you require this information in another language, large print, audio (CD or tape) or braille, please e-mail the Patient Information team at patient.information@ulh.nhs.uk